

REFERENCES

- ¹ Gauderer MW, Ponsky JL, Izant RJ Jr. *Gastrostomy without laparotomy: a percutaneous endoscopic technique*. J Paediatric Surgery 1980; 15(6):872-5
- ² Stroud M, Duncan H, Nightingale J. *Guidelines for enteral feeding in adult hospital patients*. Gut 2003; 52(suppl VII):vii1-vii12
- ³ Mitchell SL, Tetroe JM. *Survival after percutaneous endoscopic gastrostomy placement in older persons*. J Gerontol A Biol Sci Med Sci 2000; 55(12):M735-9
- ⁴ Grant MD, Rudberg MA, Brody JA. *Gastrostomy placement and mortality among hospitalized Medicare beneficiaries*. JAMA 1998; 279(24):1973-76
- ⁵ Marik PE. *Primary care: aspiration pneumonitis and aspiration pneumonia*. N Engl J Med 2001; 344(9):665-71
- ⁶ Finucane TE, Christmas C, Travis K. *Tube feeding in patients with advanced dementia: A review of the evidence*. JAMA 1999; 282(14):1365-70
- ⁷ Murphy LM, Lipman TO. *Percutaneous endoscopy does not prolong survival in patients with dementia*. Arch Intern Med 2003;163(11):1351-53
- ⁸ Saunders DS, Carter MJ, D'Silva J, James G, Bolton RP, Bardhan KD. *Survival analysis in percutaneous endoscopic gastrostomy feeding: a worse outcome in patients with dementia*. Am Journal Gastroenterol 2000; 95(6):1472-5
- ⁹ Nair S, Hertan H, Pitchumoni CS. *Hypoalbuminaemia is a poor predictor of survival after percutaneous endoscopic gastrostomy in elderly patients with dementia*. Am J Gastroenterol 2000; 95(1):133-6
- ¹⁰ Billings JA. *Recent advances: palliative care*. BMJ 2000; 321(7260):555-8
- ¹¹ *Withholding and withdrawing life-prolonging treatments: good practice in decision-making*. General Medical Council, 2002. www.gmc-uk.org/standards/whwd.htm
- ¹² Bath PMW, Bath FJ, Smithard DG. *Interventions for dysphagia in acute stroke*. In: The Cochrane Library, Issue 3, 2004. Chichester, UK: John Wiley & Sons, Ltd.
- ¹³ Norton B, Homer-Ward M, Donnelly MT, Long RG, Holmes GK. *A randomised prospective comparison of percutaneous endoscopic gastrostomy and nasogastric feeding after acute dysphagic stroke*. BMJ 1996, 312(7022):13-16

- ¹⁴ *Antibiotic prophylaxis in gastrointestinal endoscopy*. BSG guidelines in gastroenterology, British Society of Gastroenterology, 2001. www.bsg.org.uk/pdf_word_docs/prophylaxis2001.pdf
- ¹⁵ Ahmad I, Mouncher A, Abdoolah A, Stenson R, Wright J, Daniels A, Tillett J, Hawthorne AB, Thomas G. *Antibiotic prophylaxis for percutaneous gastrostomy - a prospective, randomised, double-blind trial*. *Aliment Pharmacol Ther* 2003; 18(2):209-15
- ¹⁶ Prelik G, Grune S, Leser HG, Lebherz J, Heldwein W, Machka K, Holstege A, Kern WV. *Prospective, randomised, double-blind trial of prophylaxis with single dose co-amoxiclav before percutaneous endoscopic gastrostomy*. *BMJ* 1999; 319(7214):881-4
- ¹⁷ Saunders S, Carter MJ. *Antibiotic prophylaxis after percutaneous endoscopic gastrostomy insertion: All encompassing study is needed (letter)*. *BMJ* 2000; 320(7238):870