

REFERENCES

¹ *Implementing and ensuring safe sedation practice for healthcare procedures in adults.* UK Academy of Medical Royal Colleges. London, 2001. www.rcoa.ac.uk/docs/safesedationpractice.pdf

² *Safety and sedation during endoscopic procedures.* British Society of Gastroenterology, 2003. www.bsg.org.uk/clinical_prac/guidelines/sedation.htm

³ Bowles CJ, Leicester R, Romaya C, Swarbrick E, Williams CB, Epstein O. *A prospective study of colonoscopy practice in the UK today: are we adequately prepared for national colorectal cancer screening tomorrow?* Gut 2004; 53(2):277-283

⁴ Quine MA, Bell GD, McCloy RF, Charlton JE, Develin HB, Hopkins A. *Prospective audit of upper gastrointestinal endoscopy in two regions of England: safety, staffing and sedation methods.* Gut 1995; 36(3):462-467

⁵ Whitwam JG. *Flumazenil and midazolam in anaesthesia.* Acta Anaesthesiol Scand Suppl 1995; 108:15-22

⁶ *Provision of endoscopy related services in District General Hospitals.* BSG Working Party Report, 2001. www.bsg.org.uk/pdf_word_docs/endo_related_services.pdf

⁷ Schenck J, Muller CH, Lubbers H, Mahlke R, Lehnick D, Lankisch PG. *Does gastroscopy induce myocardial ischemia in patients with coronary heart disease?* Endoscopy 2000; 32(5):373-6

⁸ Seinela L, Reinikainen P, Ahvenainen J. *Effect of upper gastrointestinal endoscopy on cardiopulmonary changes in very old patients.* Arch Gerontol Geriatr 2003; 37(1):25-32

⁹ Wang CY, Ling LC, Cardosa MS, Wong AK, Wong NW. *Hypoxia during upper gastrointestinal endoscopy with and without sedation and the effect of pre-oxygenation on oxygen saturation.* Anaesthesia 2000; 55(7): 654-8

¹⁰ Alcain G, Guillen P, Escolar A, Moreno M, Martin L. *Predictive factors of oxygen desaturation during upper gastrointestinal endoscopy in nonsedated patients.* Gastrointest Endosc 1998; 48(2):143-7

¹¹ Oei-Lim VL, Kalkman CJ, Bartelsman JF, Res JC, van Wezel HB. *Cardiovascular responses, arterial oxygen saturation and plasma catecholamine concentration during upper gastrointestinal endoscopy using conscious sedation with midazolam or propofol.* Eur J Anaesthesiol 1998; 15(5):535-43

¹² Yen D, Hu SC, Chen LS, Liu K, Kao WF, Tsai J, Chern CH, Lee CH. *Arterial oxygen desaturation during emergent nonsedated upper gastrointestinal endoscopy in the emergency department*. Am J Emerg Med 1997; 15(7):644-7

¹³ Holm C, Christensen M, Rasmussen V, Schulze S, Rosenberg J. *Hypoxaemia and myocardial ischaemia during colonoscopy*. Scand J Gastroenterol 1998; 33(7):769-72

¹⁴ Bell GD, McCloy RF, Charlton JE, Campbell D, Dent NA, Gear MW, Logan RF, Swan CH. *Recommendations for standards of sedation and patient monitoring during gastrointestinal endoscopy*. Gut 1991. 32(7): 823-7